

NEW BOUNDS ON R-PARITY VIOLATING COUPLINGS

MARC SHER

*Physics Department, College of William and Mary,
Williamsburg, VA 23187 USA*

Though the minimal supersymmetric standard model (MSSM) ¹ is a leading candidate for new physics beyond the standard model, the conservation of R-parity, R_p , which is assumed in the model has no real theoretical justification. This has motivated many authors ² to consider alternatives in which R_p is explicitly broken. In such models, sparticles can decay into non-supersymmetric particles alone, leading to novel signatures in search experiments and unusual decay processes.

The most general R_p -violating superpotential that one can write with the MSSM superfields, in the usual notation, is

$$W = \lambda_{ijk} L_i L_j \bar{E}_k + \lambda'_{ijk} L_i Q_j \bar{D}_k + \lambda''_{ijk} \bar{U}_i \bar{D}_j \bar{D}_k \quad (1)$$

Here, i, j, k are generation indices and we have rotated away a term of the form $\mu_{ij} L_i H_j$. Since the λ_{ijk} term is symmetric under exchange of i and j , and antisymmetric in color, it must be antisymmetric in flavor, thus we have $\lambda_{ijk} = -\lambda_{jik}$. Similarly, $\lambda'_{ijk} = -\lambda_{ikj}$. The number of couplings is then 36 lepton nonconserving couplings (9 of the λ type and 27 of the λ' type) and 9 baryon nonconserving couplings (all of the λ'' type) in total.

It is generally thought that λ, λ' type couplings cannot coexist with λ'' type couplings since both baryon and lepton number violations would lead to too rapid a proton decay. For this reason, previous authors have considered either lepton nonconserving *or* baryon nonconserving couplings, but not both. We will first make this assumption and consider the baryon number violating λ'' couplings alone, since most of the earlier effort has been focused on λ and λ' couplings ³. Later, we will examine proton decay in the presence of all three types of couplings.

We can write the 9 different λ'' couplings as $\lambda''_{tbs}, \lambda''_{tbd}, \lambda''_{tsd}, \lambda''_{cbs}, \lambda''_{cbd}, \lambda''_{csd}, \lambda''_{ubs}, \lambda''_{ubd}$ and λ''_{usd} . Let us first recount the existing constraints on these couplings. Brahmachari and Roy ⁴ showed that the requirement of perturbative unification typically places a bound of between 1.10 and 1.25 on many of the couplings. This was generalized to all the couplings by Goity and Sher ⁵. The latter also showed, following earlier work ^{6,7}, that $|\lambda''_{usd}|$ can be strongly bounded by the nonobservation of double nucleon decay into two kaons (such as $^{16}O \rightarrow ^{14}C K^+ K^+$, which would have been seen

in water Cerenkov detectors), and $|\lambda''_{ubd}|$ can be strongly bounded by the nonobservation of neutron-antineutron oscillations. Their bounds, for squark masses of 300 GeV, were $|\lambda''_{ubd}| < 5 \times 10^{-3}$ and $|\lambda''_{usd}| < 10^{-6}$. In the work of Barbieri and Masiero ⁷, some bounds on products of couplings were obtained by considering $K-\bar{K}$ mixing; these bounds will be discussed shortly. Finally, bounds from vertex corrections to the decay of the Z into two charged leptons and into $\bar{b}b$ have recently been obtained ⁸; though potentially interesting, with present data they are not significantly better than the bound from perturbative unification.

In this Talk, I note that many additional and interesting bounds on the λ'' couplings can be obtained by considering rare two-body nonleptonic decays of B mesons. Let us begin by considering the implications for λ'' couplings from such processes. Since lepton number is assumed to be conserved, only $\Delta B = 0$ and $\Delta B = 2$ decays can occur. Thus any bounds will be on the products of two λ'' couplings. Furthermore, since any B-decay will change the number of "b-flavors" by one unit, bounds from there will apply to products of the form $\lambda_{u_i b s} \lambda_{u_j s d}$ or $\lambda_{u_i b d} \lambda_{u_j s d}$. We first consider $\Delta B = 0$ (baryon number conserving) decays which actually give the best bounds and later comment on the $\Delta B = 2$ processes.

The details of the calculation can be found in the paper of Carlson, Roy and Sher ⁹. There, it is shown that the best bounds come from the decay $B^+ \rightarrow \bar{K}^0 K^+$. This has an extremely small rate in the Standard Model, being penguin-suppressed and also reduced by the small CKM element V_{ub} in the amplitude. The dominant diagrams contributing to this process for nonzero λ'' couplings come from squark exchange. We find that

$$\text{B.R.}(B^+ \rightarrow \bar{K}^0 K^+) \simeq 1.97 |\lambda''_{tbs} \lambda''_{tsd}|^2 (m_W/m_t)^4. \quad (2)$$

On using the recent experimental upper bound ¹⁰ of 5×10^{-5} on the branching ratio, and noting that one can replace the scalar top with a scalar charm or scalar up, we then have

$$\frac{|\lambda''_{qbs} \lambda''_{qsd}| m_W^2}{m_q^2} < 5 \times 10^{-3}, \quad (3)$$

for $q = t, c, u$.

One can repeat the calculation for the decay $B^+ \rightarrow \bar{K}^0 \pi^+$ (or equivalently $B^- \rightarrow K^0 \pi^-$) in much the same way. The result (with only the t-squark contribution being considered) is

$$\frac{|\lambda''_{qbd} \lambda''_{qsd}| m_W^2}{m_q^2} < 4.1 \times 10^{-3}, \quad (4)$$

for $q = t, c, u$.

Additional bounds were obtained by Barbieri and Masiero ⁷ from the contribution of $K-\bar{K}$ mixing to the K_L-K_S mass difference. We have generalized their results by including effects from all of the squarks, updated top mass and CKM angles, etc., and find bounds on several combinations of couplings (including the product $|\lambda''_{tbs} \lambda''_{tbd}|$), the plot can be found in Ref. [9].

What about $\Delta B = 2$ decays? One can envision the decay $B \rightarrow \Sigma^+ \Sigma^-$ or $\Lambda \Lambda$. However, a simple estimate of the rate gives branching ratios (assuming the B-violating couplings are unity and the scalar quark masses are near the W mass) of $O(10^{-8})$. The smallness of the rate is due in large part to two small CKM elements in the amplitude. Thus, such processes will not provide interesting bounds unless 10^{10} B-decays are studied.

It is generally assumed that the presence of both lepton nonconserving terms and baryon nonconserving terms leads to unacceptably rapid proton decay. However, if enough third generation fields are involved, proton decay can be sufficiently suppressed as to make some of the bounds very weak (or, in a few cases, nonexistent). To see this, suppose both λ and λ'' terms exist. Consider the bound on the product $|\lambda_{\mu\tau\tau} \lambda''_{usd}|$. Although there is a suppression due to mixing angles and heavy squark propagators, the proton lifetime bound gives a strong bound of $|\lambda_{\mu\tau\tau} \lambda''_{usd}| < 10^{-14}$. This bound is independent of the final state leptons, and thus applies to all 9 of the λ couplings. Similar bounds can be obtained for all λ'' couplings with at most one heavy field (which is then the scalar quark); we obtain $|\lambda_{ijk} \lambda''_{ubd}| < 10^{-13}$, $|\lambda_{ijk} \lambda''_{ubs}| < 10^{-12}$, $|\lambda_{ijk} \lambda''_{csd}| < 10^{-13}$ and $|\lambda_{ijk} \lambda''_{tsd}| < 10^{-12}$. However, if the λ'' coupling has two heavy fields, a loop is necessary. This gives much weaker bounds; we obtain $|\lambda_{ijk} \lambda''_{tbs}| < 10^{-2}$, $|\lambda_{ijk} \lambda''_{tbd}| < 10^{-3}$, $|\lambda_{ijk} \lambda''_{cbs}| < 10^{-3}$ and $|\lambda_{ijk} \lambda''_{cbd}| < 10^{-2}$. We thus see that the lack of observation of proton decay does NOT always give very strong bounds on the product of the lepton number violating and baryon number violating couplings.

Finally, we consider the product of λ' and λ'' couplings. Here, there are 27×9 possible products, of the form $|\lambda'_{ijk} \lambda''_{abc}|$. We have examined all possible products of couplings and found that the vast majority are tightly bounded (product is less than 10^{-6}), but some are not. Rather than list the bounds for all 243 combinations, only the bounds which are greater than 10^{-6} (for the product of the λ' and λ'' couplings) will be

given explicitly. It is found that all products with a λ'_{usd} , λ''_{ubd} and λ''_{ubs} are smaller than 10^{-9} . The same is true for λ''_{csd} , λ''_{cbd} and λ''_{cbs} , *except* for $|\lambda'_{tbl} \lambda''_{csd}|$ which is $< 10^{-1}$, $|\lambda'_{ust} \lambda''_{cbd}|$ which is $< 10^{-2}$ and $|\lambda'_{udl} \lambda''_{cbs}|$, for which no bound better than the unitarity bound could be found. Here l is any lepton. For λ''_{tsd} , the only bound which is not very small is the combination $|\lambda'_{cbl} \lambda''_{tsd}|$, which is bounded only by unitarity. For λ''_{tbd} and λ''_{tbs} , we find $|\lambda'_{ud(e,\mu)} \lambda''_{tbd}| < 10^{-2}$, $|\lambda'_{us(e,\mu)} \lambda''_{tbs}| < 10^{-2}$, $|\lambda'_{c(d,s)(e,\mu)} \lambda''_{tbd}| < 10^{-3}$, $|\lambda'_{c(d,s)(e,\mu)} \lambda''_{tbs}| < 10^{-3}$, $|\lambda'_{cd\tau} \lambda''_{tbd}| < 10^{-5}$, $|\lambda'_{cs\tau} \lambda''_{tbs}| < 10^{-5}$, whereas $|\lambda'_{(u,c)d\tau} \lambda''_{tbs}|$ and $|\lambda'_{(u,c)s\tau} \lambda''_{tbd}|$ are bounded only by unitarity. All other bounds are quite tiny. Thus, of the 243 combinations of couplings, thirty have bounds greater than 10^{-6} , and eight are completely unconstrained by the lack of observation so far of proton decay.

It is interesting that the standard supersymmetric model can contain some baryon number and lepton number violating coupling constants which are of order unity, and which do not lead to excessively fast proton decay. Such couplings could be measured when squarks and sleptons are discovered, since they will lead to baryon and lepton number violating squark and slepton decays. These have been extensively discussed by Dreiner and Ross and by Dimopoulos, et al. ¹¹, who analyze the impact of such decays on phenomenology.

1. H. P. Nilles, Phys. Rep. **110** (1984) 1; H.E. Haber and G.L. Kane, Phys. Rep. **117** (1985) 95.
2. J. Ellis, et al., Phys. Lett. **B150** (1985) 142; G.G. Ross and J.W.F. Valle, Phys. Lett. **B151** (1985) 375; S. Dawson, Nucl. Phys. **B261** (1985) 297; S. Dimopoulos and L. Hall, Phys. Lett. **B207** (1987) 210.
3. L. Hall and M. Suzuki, Nucl. Phys. **B231** (1984) 419; V. Barger, G.F. Giudice and T. Han, Phys. Rev. **D40** (1989) 2987; R. Godbole, P. Roy and X. Tata, Nucl. Phys. **B401** (1993) 67.
4. B. Brahmachari and P. Roy, Phys. Rev. **D50** (1994) R39.
5. J.L. Goity and M. Sher, Phys. Lett. **B346** (1995) 69.
6. F. Zwirner, Phys. Lett. **B132** (1983) 103.
7. R. Barbieri and A. Masiero, Nucl. Phys. **B267** (1986) 679.
8. G. Bhattacharyya, D. Choudhury and K. Sridhar, CERN preprint CERN-TH-95-89, hep-ph 9504314.
9. C.E. Carlson, P. Roy and M. Sher, Physics Letters, in press (1995).
10. F. Würthwein, CLEO Collaboration, talk given at the meeting of the American Physical Society, Washington DC, April, 1995; Cornell Preprint No. CLNS 95/1338, in preparation.).
11. H. Dreiner and G.G. Ross, Nucl. Phys. **B365** (1991) 597; S. Dimopoulos, R. Esmailzadeh, L. Hall,

J.-P. Merlo and G. Starkman, Phys. Rev. **D41**
(1990) 2099.